

N-JORDAN HOMOMORPHISMS

M. ESHAGHI GORDJI

ABSTRACT. Let $n \in \mathbb{N}$, and let A, B be two rings. An additive map $h : A \rightarrow B$ is called n -Jordan homomorphism if $h(a^n) = (h(a))^n$ for all $a \in A$. Every Jordan homomorphism is an n -Jordan homomorphism, for all $n \geq 2$, but the converse is false, in general. In this paper we investigate the n -Jordan homomorphisms on Banach algebras. Indeed some results related to continuity are given as well.

1. INTRODUCTION AND PRELIMINARIES

Let A, B be two rings (algebras). An additive map $h : A \rightarrow B$ is called n -Jordan homomorphism (n -ring homomorphism) if $h(a^n) = (h(a))^n$ for all $a \in A$, ($h(\prod_{i=1}^n a_i) = \prod_{i=1}^n h(a_i)$, for all $a_1, a_2, \dots, a_n \in A$). If $h : A \rightarrow B$ is a linear n -ring homomorphism, we say that h is n -homomorphism. The concept of n -homomorphisms was studied for complex algebras by Hejazian, Mirzavaziri, and Moslehian [3] (see also [1] and [7]). A 2-Jordan homomorphism is a Jordan homomorphism, in the usual sense, between rings. Every Jordan homomorphism is an n -Jordan homomorphism, for all $n \geq 2$, (see for example Lemma 6.3.2 of [6]), but the converse is false, in general. For instance, let A be an algebra over $\mathbb{C}$ and let $h : A \rightarrow A$ be a non-zero Jordan homomorphism on A . Then $-h$ is a 3-Jordan homomorphism. It is easy to check that $-h$ is not 2-Jordan homomorphism or 4-Jordan homomorphism. The study of ring homomorphisms between Banach algebras A and B is of interest even if $A = B = \mathbb{C}$. For example the zero mapping, the identity and the complex conjugate are ring homomorphisms on $\mathbb{C}$, which are all continuous. On the other hand the existence of a discontinuous ring homomorphism on $\mathbb{C}$ is well-known. More explicitly, if G is the set of all surjective ring homomorphisms on $\mathbb{C}$, then $Card(G) = 2^{Card(\mathbb{C})}$. In fact, Charnow [2; Theorem 3] proved that there exist $2^{Card(\mathbb{C})}$ automorphisms for every algebraically closed field K . It is also known that if A is a uniform algebra on a compact metric space, then there are exactly $2^{Card(\mathbb{C})}$ complex-valued ring homomorphisms on A whose kernels are non-maximal prime ideals (see [4;

2000 *Mathematics Subject Classification.* Primary 47B48; Secondary 46L05, 46H25.

Key words and phrases. Jordan homomorphism; n -homomorphism; Banach algebra.

Corollary 2.4]). As an example, take

$$\mathcal{A} := \begin{bmatrix} 0 & \mathbb{R} & \mathbb{R} & \mathbb{R} \\ 0 & 0 & \mathbb{R} & \mathbb{R} \\ 0 & 0 & 0 & \mathbb{R} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

then $\mathcal{A}$ is an algebra equipped with the usual matrix-like operations. It is easy to see that

$$\mathcal{A}^3 \neq 0 = \mathcal{A}^4.$$

So any additive map from $\mathcal{A}$ into itself is a 4-Jordan homomorphism, but its kernel does not need to be an ideal of $\mathcal{A}$. Let now $\mathcal{B}$ be the algebra of all $\mathcal{A}$ -valued continuous functions from $[0, 1]$ into $\mathcal{A}$ with sup-norm. Then $\mathcal{B}$ is an infinite dimension Banach algebra, also product of any four elements of $\mathcal{B}$ is 0. Since $\mathcal{B}$ is infinite dimension, there are linear discontinuous maps which are 4-Jordan homomorphisms from $\mathcal{B}$ into itself (see [3]). In this paper we study the continuity of linear n -Jordan homomorphisms on C^* -algebras.

2. MAIN RESULT

By definition, it is obvious that n -ring homomorphisms are n -Jordan homomorphisms. Conversely, under a certain condition, n -Jordan homomorphisms are ring homomorphisms. For example, each Jordan homomorphism h from a commutative Banach algebra A into $\mathbb{C}$ is a ring homomorphism: Fix $a, b \in A$ arbitrarily. Since $h((a+b)^2) = (h(a+b))^2$, a simple calculation shows that $h(ab+ba) = 2h(a)h(b)$. The commutativity of A implies $h(ab) = h(a)h(b)$, and hence h is a ring homomorphism. In 1968, Zelazko [8] proved the following Theorem (see also Theorem 1.1 of [5]).

Theorem 2.1. *Suppose A is a Banach algebra, which need not to be commutative, and suppose B is a semisimple commutative Banach algebra. Then each Jordan homomorphism $h : A \rightarrow B$ is a ring homomorphism.*

We prove the following result for 3-Jordan homomorphisms and 4-Jordan homomorphisms on commutative algebras.

Theorem 2.2. *Let $n \in \{3, 4\}$ be fixed, A, B be two commutative algebras, and let $h : A \rightarrow B$ be a n -Jordan homomorphism. Then h is n -ring homomorphism.*

Proof. First let $n = 3$. Recall that h is additive mapping such that $h(a^3) = (h(a))^3$ for all $a \in A$. Replacement of a by $x + y$ results in

$$h(x^2y + xy^2) = h(x)^2h(y) + h(x)h(y)^2. \quad (1)$$

Hence for every $x, y, z \in A$ we have

$$\begin{aligned} h(xyz) &= \frac{1}{2}h\{(x+z)^2y + (x+z)y^2 - (x^2y + xy^2 + z^2y + zy^2)\} \\ &= \frac{1}{2}\{h[(x+z)^2y + (x+z)y^2] - h[x^2y + xy^2] - h[z^2y + zy^2]\} \\ &= \frac{1}{2}\{[h(x+z)]^2h(y) + (x+z)[h(y)]^2 - [h(x)]^2h(y) + h(x)[h(y)]^2 \\ &\quad - [h(z)]^2h(y) + h(z)[h(y)]^2\} \\ &= h(x)h(y)h(z). \end{aligned}$$

This means that h is a 3-ring homomorphism. Now suppose $n = 4$. Then h is additive and $h(a^4) = (h(a))^4$ for all $a \in A$. Replacing a by $x + y$ in above equality to get

$$h(4x^3y + 6x^2y^2 + 4xy^3) = 4h(x)^3h(y) + 6h(x)^2h(y)^2 + 4h(x)h(y)^3. \quad (2)$$

Replacing x by $x + z$ in (2), we obtain

$$\begin{aligned} h\{(4x^3y + 6x^2y^2 + 4xy^3) + (4z^3y + 6z^2y^2 + 4zy^3) + 12(x^2zy + xz^2y + xzy^2)\} \\ = (4h(x)^3h(y) + 6h(x)^2h(y)^2 + 4h(x)h(y)^3) + (4h(z)^3h(y) + 6h(z)^2h(y)^2 \\ + 4h(z)h(y)^3) + 12(h(x)^2h(z)h(y) + h(x)h(z)^2h(y) + h(x)h(z)h(y)^2). \end{aligned} \quad (3)$$

Combining (2) by (3) to get

$$h\{(xyz)(x + y + z)\} = (h(x)h(y)h(z))(h(x) + h(y) + h(z)). \quad (4)$$

Replacing z by $-x$ in (4) to obtain

$$h(x^2y^2) = h(x)^2h(y)^2 \quad (5)$$

replacing y by $y + w$ in (5), we get

$$h(x^2yw) = h(x)^2h(y)h(w). \quad (6)$$

Now replace x by $x + t$ to obtain

$$h(xtyw) = h(x)h(t)h(y)h(w)$$

hence, h is 4-ring homomorphism.

□

By Theorem 2.2 and Theorem 3.2 of [1], we conclude the following result.

Corollary 2.3. *Let $h : A \rightarrow B$ be a linear involution preserving 3-Jordan homomorphism between commutative C^* -algebras. Then h is norm contractive ($\|h\| \leq 1$).*

Also by above Theorem and Theorem 2.3 of [7], we have the following.

Corollary 2.4. *Let $h : A \rightarrow B$ be a linear involution preserving 4-Jordan homomorphism between commutative C^* -algebras, then h is completely positive. Thus h is bounded.*

Now we prove our main Theorem.

Theorem 2.5. *Suppose A is a Banach algebra, which need not to be commutative, and suppose B is a semisimple commutative Banach algebra. Then each 3-Jordan homomorphism $h : A \rightarrow B$ is a 3-ring homomorphism.*

Proof. We prove the Theorem in two steps as follows.

STEP I. Suppose $B = \mathbb{C}$. We have $h(a^3) = h(a)^3$ for all $a \in A$. Replace a by $x + y$ to obtain

$$h(xyx + yx^2 + y^2x + x^2y + xy^2 + yxy) = 3(h(x)^2h(y) + h(x)h(y)^2) \quad (7)$$

replace y by $-y$ in (7) to get

$$h(-xyx - yx^2 + y^2x - x^2y + xy^2 + yxy) = 3(-h(x)^2h(y) + h(x)h(y)^2). \quad (8)$$

By (7), (8), we obtain the relation

$$h(xy^2 + y^2x + yxy) = 3(h(x)h(y)^2). \quad (9)$$

Replacing y by $y - z$ in (9), we get

$$\begin{aligned} h(xy^2 + xz^2 - 2xyz + yxy - yxz - zxy + zxz + z^2x + y^2x - 2yzx) \\ = 3(h(x)^2h(y) + h(x)h(y)^2) - 6h(x)h(y)h(z). \end{aligned} \quad (10)$$

By (9) and (10), we obtain

$$h(yxz + zxy + 2xyz + 2yzx) = 6h(x)h(y)h(z) \quad (11)$$

replacing z by x in (11) to get

$$h(3yx^2 + x^2y + 2xyx) = 6h(x)^2h(y) \quad (12)$$

combining (9) and (12) to obtain

$$h(xyx + 2yx^2) = 3h(x)^2h(y). \quad (13)$$

By (8) and (13), we conclude that

$$h(yx^2 - x^2y) = 0. \quad (14)$$

Replacing x by $x + z$ in (14) to get

$$h(yx^2 + yz^2 + 2yxz - x^2y - z^2y - 2xzy) = 0$$

by above equality and (14) it follows that

$$h(yxz - xzy) = 0. \quad (15)$$

Combining (11) and (15), we obtain

$$h(yxz + 3xyz + 2yzx) = 6h(x)h(y)h(z) \quad (16)$$

replace z by x in (16), we get

$$h(xyx + yx^2) = 2h(x)^2h(y) \quad (17)$$

combining (13) and (17) to obtain

$$h(yx^2) = h(y)h(x)^2 \quad (18)$$

replace x by $x + z$ in (18), we conclude that

$$h(yxz) = h(y)h(x)h(z)$$

hence h is 3-ring homomorphism.

STEP II. B is arbitrary semisimple and commutative. Let M_B be the maximal ideal space of B . We associate to each $f \in M_B$ a function $h_f : A \rightarrow \mathbb{C}$ defined by

$$h_f(a) := f(h(a))$$

for all $a \in A$. It is easy to see that h_f is additive and $h_f(a^3) = (h_f(a))^3$ for all $a \in A$. So STEP I applied to h_f , implies that h_f is a 3-ring homomorphism. By the definition of h_f , we obtain that

$$f(h(abc)) = f(h(a))f(h(b))f(h(c)) = f(h(a)h(b)h(c)).$$

Hence

$$h(abc) - h(a)h(b)h(c) \in \text{Ker}(f)$$

for all $a, b, c \in A$ and all $f \in M_B$. Since B is assumed to be semisimple, we get $h(abc) = h(a)h(b)h(c)$ for all $a, b, c \in A$. We thus conclude that h is a 3-ring homomorphism, and the proof is complete. $\square$

From now on we consider such n -Jordan homomorphisms that are linear.

Corollary 2.6. *Suppose A, B are C^* -algebras, which A need not to be commutative, and suppose B is semisimple and commutative. Then every involution preserving 3-Jordan homomorphism $h : A \rightarrow B$ is norm contractive ($\|h\| \leq 1$).*

Proof. It follows from Theorem 2.3 above and Theorem 2.1 of [1]. $\square$

Theorem 2.7. *Let $h : A \rightarrow B$ be a bounded involution preserving k -Jordan homomorphism between C^* -algebras such that $h(a^*a) = h(a)^*h(a)$ for all $a \in A$. Then h is norm contractive ($\|h\| \leq 1$).*

Proof. By using Lemma 2.4 of [7], we have

$$\begin{aligned}
\|h(a)\|^{4k+2} &= \|(h(a)^*h(a))^{2k+1}\| = \|(h(a)^*h(a))^k(h(a)^*h(a))(h(a)^*h(a))^k\| \\
&= \|[h(a)(h(a)^*h(a))^k]^*[h(a)(h(a)^*h(a))^k]\| \\
&= \|h(a)(h(a)^*h(a))^k\|^2 = \|h(a)(h(a^*a))^k\|^2 \\
&= \|h(a)(h(a^*a)^k)\|^2 \leq \|h(a)\|^2\|h((a^*a)^k)\|^2 \\
&\leq \|h\|^2\|a\|^2\|h\|^2\|(a^*a)^k\|^2 \\
&\leq \|h\|^4\|a\|^{4k+2},
\end{aligned}$$

for all $a \in A$. Which implies that $\|h\| \leq 1$ by taking $4k + 2$ -th roots. $\square$

REFERENCES

- [1] J. Bracic and M. S. Moslehian, *On automatic continuity of 3-homomorphisms on Banach algebras*, Bull. Malays. Math. Sci. Soc. (2) 30 (2007), no. 2, 195–200.
- [2] Charnow, A. The automorphisms of an algebraically closed field. Canad. Math. Bull. 13 1970 95–97.
- [3] Sh. Hejazian, M. Mirzavaziri and M. S. Moslehian, *n -homomorphisms*, Bull. Iranian Math. Soc. 31 (2005), no. 1, 13–23, 88. (Reviewer: Peter emrl) 47B48 (16N60 46H05)
- [4] T. Miura, S.-E. Takahasi and N. Niwa, Prime ideals and complex ring homomorphisms on a commutative algebra. Publ. Math. Debrecen 70 (2007), no. 3-4, 453–460. (Reviewer: Armando R. Villena) 46J10
- [5] T. Miura, S.-E. Takahasi and G. Hirasawa, Hyers-Ulam-Rassias stability of Jordan homomorphisms on Banach algebras. J. Inequal. Appl. 2005, no. 4, 435–441.
- [6] T. Palmer, Banach algebras and the general theory *-algebras, Vol I. Cambridge: Univ Press (1994).
- [7] E. Park and J. Trout, *On the nonexistence of nontrivial involutive n -homomorphisms of C^* -algebras*, To appear in Trans. Amer. Math. Soc.
- [8] Zelazko, W. A characterization of multiplicative linear functionals in complex Banach algebras. Studia Math. 30 1968 83–85. (Reviewer: B. Yood) 46.50

DEPARTMENT OF MATHEMATICS, SEMNAN UNIVERSITY, P. O. BOX 35195-363, SEMNAN, IRAN

E-mail address: madjid.eshaghi@gmail.com